INDEX

Asker, A. A., *see* Robertson, A.

Barlow, H. W. B., *see* Jewitt, T. N.
Black, W., breeding for disease-resistance in potatoes, 116–24.
B-vitamins in grass, 170–8; experimental technique, 172–3; grasses studied, 171; manurial treatments and their effects, 172, 176; vitamin-contents of grasses from different plots, 173–5.

Cabbages, ascorbic acid in, 28–36; distribution in leaf and head, 28–9; diurnal variation, 30; effect of maturity, 30–1; of time of picking, 31–2; of variety, size of head, and solar radiation, 32–4; methods of analysis, 28, and sampling, 29–30.
Campbell, G. K. G., *see* Hill, A. G. G.
Cattle, dairy, pedigree, movement of two breeds of, 125–31; 'county' replacements of Friesians and Shorthorns, 127–9; measurement of migration, 125–7.
Cereals, breeding for disease-resistance in Canada. Pt. I. Cereals, 133–40; barley, 138–40; oats, 136–8; wheat, 134–6. Pt. II. Flax and forage crops, 222–8; alfalfa (lucerne), 225–7; flax, 222–4; forage crops, 224–8; legumes, 224–5; red clover, 227–8.
Composts incorporating human faeces, field tests with, in N. China, 73–82; experimental technique, 73–4, 75–6, 77; moisture and nitrate changes, 78–80; nutrients removed by crops, 77–8; summaries of results, 74–5, 76–7, 80–1.
Corkill, I., pasture improvement in New Zealand, 157–69.
Cotton, planting date and production of, in Buganda province of Uganda, 245–58; meteorology, 245–8; multiplication scheme, 252–3; provincial production, 254–6; soil nitrogen, 253–4; sowing-date experiment, 248–50; variety trials, 250–2.

Dairy herd fertility, 229–37; conception rate, 230–3; N.I.R.D. herd, 230; potency of bulls, 233–4; pregnancy diagnosis, 235; proportion of effective matings, 233; veterinary treatment, 234–5.
Digestibility studies, application of modified procedures in, 188–92; analytical methods used, 189; data of food and faeces (rabbits), 190; recovery of lignin, silica, iron as reference substances, 189–91.
Disease-resistance (in plants), breeding for: cereals in Canada, 133–40; flax and forage crops in Canada, 222–8; oats, 199–204, 205–14; potatoes, 116–24, 238–44; wheat in S. Australia, 193–8.
Druce, E., application of modified procedures in digestibility studies, 188–92.
Duckworth, J., the three-quarter-bred Holstein-Zebu heifer. Pt. III. The age of puberty, 23–7.

Eelworm, the potato-root, in the U.K., 60–71; areas affected, 60–2; attack on potatoes and tomatoes, 62–3; control methods, 66–8; cropping infested land, 69–71; field surveys, 61, 68–9; how infestation spreads, 64–5; 'potato sickness', 63–4; soil types, influence of, 65–6.
Ellison, W., hill-land reclamation in Montgomeryshire, Wales, 215–21.
Excreta, animal, in the Sudan Gezira, 1–17; dung from field collections and penned animals, 2–3; field experiments, 12–14; nitrification experiments, 7–8; nitrogen in dung and urine, 3–6; pot experiments, 8–12.

Flax and forage crops, breeding for disease-resistance in Canada, 222–8; alfalfa and bacterial wilt, 225–6, and blackstem, 227, and winter crown rot, 226–7; crown rot in red clover, 229–30; diseases in legumes, 224–5; flax, 222–4; forage crops, 224.
Foot, A. S., dairy herd fertility, 229–37.
Fruit-trees, prolonged dormancy of deciduous, in warm climates, 259–64; remedies: orchard management, 262, selection and breeding, 261–2, spray treatment, 262–3; the phenomenon, 259–60.

Goulden, C. H., breeding for disease-resistance in Canada. Pt. I. Cereals, 133–40. Pt. II. Flax and forage crops, 222–8.

Hallsworth, E. G., ascorbic acid in cabbages, 28–36.
Heifer, the three-quarter-bred Holstein-Zebu. Pt. III. The age of puberty, 23–7. Relation between age, weight, and puberty, 23–5; between age of puberty and growth-rate, 25–6.
Hill, A. G. G., prolonged dormancy of deciduous fruit-trees in warm climates, 259–64.
Hill-land reclamation in Montgomeryshire, Wales, 215–21; areas and soils, 215–16; establishment of clovers, 219–20; initial cultivation, 216–17; pioneer crops, 218–19; potatoes, 217–18; problems, 215; re-seeding, 219; sugar-beet, 218.
Hopkins, E. S., wheat production in Canada, 92–104.

Jewitt, T. N., animal excreta in the Sudan Gezira, 1–17.
Jones, E. T., breeding of disease-resistant varieties of oats. Pt. I. Physiologic races, diseases, and pests, 199–204. Pt. II. Inheritance of disease-resistance, 205–14.

Khishin, A. F. E., twenty years of Shorthorn breeding in Egypt, 83–92.

Land utilization in the wet tropics, 148–56; land use in upland regions, 150–2, on illuvial lands, 152–5; organic matter and nitrogen, 149–50.
Lewis, V. M., *see* Hallsworth, E. G.
Liming materials, the effectiveness of, 265–76; crop analyses, 268–9, 274–6; field technique, 266–7; liming materials used, 266, 273; soil data, 269–70, 276; soils, 265; yield data, 267–8, 273–6.
Livestock, Iraqi, 48–59; buffaloes, 54–7; cattle, 57–8; goats, 58–9; sheep, 49–54; sheep products: butter, 52, 53, mutton, 52, 53, wool, 53–4.

Manning, H. L., planting date and cotton production in the Buganda province of Uganda, 245–58.
Maule, J. P., breeding for milk in a Mediterranean environment. Pt. II. Goats, 37–47.
Milk, goats', breeding for in a Mediterranean environment, 37–47; breeding and management of goats at Government stock farm, 40; goat industry of Cyprus, 37–40, 46–7; milk yields, 39–46.

Oats, breeding of disease-resistant varieties, 199–204. Pt. I. Physiologic races, diseases, and pests, 199–204. Crown rust, 200–1; halo blight, 202; leaf-spot and seedling blight, 202; loose and covered smuts, 201; mildew, 201–2; parasitic diseases in Britain, 200 et sqq.; non-parasitic, 203; plant-pests, 202–3; seedling rot and root-rots, 202. Pt. II. Inheritance of disease-resistance, 205–14; breeding technique, 212–13; recent progress, 210–11; resistance to loose and covered smuts, 206–8, to stem and crown rusts, 208–10.

Pasture improvement in New Zealand, 157–69; breeding-work, 157–8; cocksfoot, 164–5; Italian rye-grass, 160–1, 162; perennial rye-grass, 158–60, 162, 163; red clover, 166–7; short-rotation rye-grass, 161–3; timothy, 165; Western Wolths rye-grass, 164; white clover, 165–6; use of pedigree strains to increase production, 167–9.
Plank, J. E. van der, relation between the size of fields and the spread of plant-disease into them. Pt. II. Diseases caused by fungi with air-borne spores, with a note on horizons of infection, 18–22. Pt. III. Examples and discussion, 141–7.
Plant-disease, relation of spread to size of fields. Pt. II. Diseases caused by fungi with air-borne spores, 18–22. Effect of enlarging fields on the multiplication of infection, 20; effect of making fields larger and fewer on the movement of spores between them, 19; horizons of infection, 20–1; scope for local hygienic measures, 22. Pt. III. Examples and discussion, 141–7. Crowd diseases and systems of land usage, 144; diseases: aphis-borne virus of beet, 141–3; rosette of ground-nuts, streak of maize, relations between size of field and eradication of sources of infection nearby, 143, and infection carried by seed or sett, 143–4.
Potatoes, breeding for disease-resistance in, 116–24; blight, 118–20; breeding methods, 122–3; leaf-roll, 121–2; mosaic, 120–1; scab, 122; wart disease, 117–18.
Potatoes, disease-resistance in, 233–44; blight, 239–42; Potato Synonym Committee, 238–9; virus disease, 242–3; wart disease, 238–9.
Pugsley, A. T., back-crossing for resistance to stem rust of wheat in South Australia, 193–8.

Reith, J. W. S., effectiveness of various liming materials, 265–76.
Ridler, B., *see* Foot, A. S.
Robertson, A., movement of two breeds of pedigree dairy cattle in Great Britain, 125–31.

Salaman, R. N., disease-resistance in potatoes: an historical note, 238–44.
Sastry, K. N. S., *see* Waite, R.
Scott, James C., field tests with composts incorporating human faeces, 73–82.
Sheep, seasonal observations on the grazing habits of, 105–15; observational technique, 105–8; distances travelled, 110, 111; mechanics of grazing, 112–14; times of grazing, resting, and idling, 108–13.
Shorthorn breeding in Egypt, twenty years of, 83–92; Egyptian Damiettas × Dairy Shorthorns, 82 sqq.: beef production, 88–9; butter-fat in milk, 87–8; diseases, 89–90; historical survey, 84–5; milk yield of crosses, 85–6, of pure Damiettas, 85, of pure Shorthorns, 86–7.
Stevenson, T. M., *see* Goulden, C. H.

Tempany, H. A., land utilization in the wet tropics, 148–56.
Thomas, Brynmor, the B-vitamins in grass, 170–8.
Thompson, H. W., the potato-root eelworm in the United Kingdom, 60–71.
Timothy (*Phleum pratense*) and some other grasses, composition of, during seasonal growth, 179–87; analytical methods used, 180; chemical composition and growth data of timothy, 180–4; correlation of protein with carotene and fibre, 182–3; grasses used, 179–80; separation of leaf and stem, 185–6.
Tribe, D. E., seasonal observations on the grazing habits of sheep, 105–15.

Vitamin-C in cabbages, *see* ascorbic acid in, 28–36.

Waite, R., composition of timothy and some other grasses during seasonal growth, 179–87.

Walker, H. F., *see* Thomas, Brynmor.
Wheat in S. Australia, back-crossing for resistance to stem rust, 193–8; back-crossing procedure, 194–6, 197–8; release of new strains, 195–6; stem rust in S. Australia, 193.
Wheat production in Canada, 92–104; areas under wheat, 93, 95–6; climates, 92, 94–5; comparisons with the U.S.A., 98; effects of hay crops in rotations, 100–1; future prospects, 104; plant-breeding achievements, 98–100; tillage practices and soil-erosion control, 103–4; use of F.Y.M. and fertilizers, 101–2; yields, 96–8, 100, 102, 104.
Willcox, J. S., *see* Druce, E.
Williams, E. G., *see* Reith, J. W. S.
Williamson, Grahame, Iraqi livestock, 48–59.

PRINTED IN
GREAT BRITAIN
AT THE
UNIVERSITY PRESS
OXFORD
BY
CHARLES BATEY
PRINTER
TO THE
UNIVERSITY